


Probing individual edge states with unprecedented precision

September 12, 2018, University of Basel



Measured tunneling current and its dependence on the two applied magnetic fields: The fans of red/yellow curves each correspond to a fingerprint of the conducting edge states. Each individual curve separately shows one of the edge states. ... more ▾

A new technique makes it possible to obtain an individual fingerprint of the current-carrying edge states occurring in novel materials such as topological insulators or 2-D materials. Physicists of the University of Basel present the new method together with American scientists in *Nature Communications*.

mediamarkt.ch
Hauptsache Ihr habt Spaß!
More information ▸

bitdefender.ch
Virenschutz & Antivirus Software
More information ▸

conrad.ch
Conrad Electronic Shop mit über 500'000 Produkten!
More information ▸

TRG AD

While insulators do not conduct electrical currents, some special materials exhibit peculiar electrical properties. Though not conducting through their bulk, their surfaces and edges may support electrical currents due to quantum mechanical effects, and do so even without causing losses. Such topological insulators have attracted great interest in recent years due to their remarkable properties. In particular, their robust edge states are very promising since they could lead to great technological

advances.

Similar effects as the edge states of such topological insulators also appear when a two-dimensional metal is exposed to a strong magnetic field at low temperatures. When the so-called quantum Hall effect is realized, current is thought to flow only at the edges, where several conducting channels are formed.

Probing individual edge states

Until now, it was not possible to address the numerous current-carrying states individually or to determine their positions separately. The new technique now makes it possible to obtain an exact fingerprint of the current-carrying edge states with nanometer resolution. This is reported by researchers of the

mediamarkt.ch
Hauptsache Ihr habt Spaß!
More information ▸


bitdefender.ch
Virenschutz & Antivirus Software
More information ▸

conrad.ch
Conrad Electronic Shop mit über 500'000 Produkten!
More information ▸

TRG AD

Featured Last comments Popular

 Gravitational waves provide dose of reality about extra dimensions 🕒 Sep 14, 2018 💬 52

 New photonic chip promises more robust quantum computers 🕒 Sep 14, 2018 💬 0

 Geologists reveal ancient connection between England and France 🕒 Sep 14, 2018 💬 7

 Gravity theory saved from death 🕒 Sep 14, 2018 💬 186

 Half-billion-year-old fossils offer new clues to how life exploded on the sea floor 🕒 Sep 14, 2018 💬 6

[more »](#)

Phys.org on Facebook

 Like 1.4M people like this. [Sign Up](#) to see what your friends like.

Email newsletter

Relevant PhysicsForums posts 

Department of Physics and the Swiss Nanoscience Institute of the University of Basel in collaboration with colleagues of the University of California, Los Angeles, as well as of Harvard and Princeton University, USA.

In order to measure the fingerprint of the conducting edge states, the physicists led by Prof. Dominik Zumbühl have further developed a technique based on tunneling spectroscopy. They have used a gallium arsenide nanowire located at the sample edge which runs in parallel to the edge states under investigation. In this configuration, electrons may jump (tunnel) back and forth between a specific edge state and the nanowire as long as the energies in both systems coincide. Using an additional magnetic field, the scientists control the momentum of tunneling electrons and can address individual edge states. From the measured tunneling currents, the position and evolution of each edge state may be obtained with nanometer precision.

Tracking the evolution

This new technique is very versatile and can also be used to study dynamically evolving systems. Upon increasing the magnetic field, the number of edge states is reduced, and their distribution is modified. For the first time, the scientists were able to watch the full edge state evolution starting from their formation at very low magnetic fields.

With increasing magnetic field, the edge states are first compressed towards the sample boundary until eventually, they move towards the inside of the sample and then disappear completely. Analytical and numerical models developed by the research team agree very well with the experimental data.

"This new technique is not only very useful to study the quantum Hall edge states," Dominik Zumbühl comments the results of the international collaboration. "It might also be employed to investigate new exotic materials such as topological insulators, graphene or other 2-D materials."

+ Explore further: Engineering electron pathways in 2-D topological insulators

More information: T. Patlatiuk et al, Evolution of the quantum Hall bulk spectrum into chiral edge states, *Nature Communications* (2018). DOI: [10.1038/s41467-018-06025-3](https://doi.org/10.1038/s41467-018-06025-3)

Journal reference: [Nature Communications](#)

Provided by: [University of Basel](#)

100 shares

feedback to editors

Is field theory needed in Regge-theory? 6 hours ago

Constant field in the Lagrangian 7 hours ago

What Determines Whether Radiation Passes Through Materials? 19 hours ago

Nature of coulomb force 19 hours ago

Higgs mechanism 20 hours ago

Nuclear and Atomic energy levels 20 hours ago

More from High Energy, Nuclear, Particle Physics


mediamarkt.ch
Hauptsache Ihr habt Spaß!
More information >

bitdefender.ch
Virenschutz & Antivirus Software
More information >

conrad.ch
Conrad Electronic Shop mit über 500'000 Produkten!
More information >

TRG AD


Related Stories



Engineering electron pathways in 2-D topological insulators 6 December 4, 2017

In a recent article published in *Physical Review Letters* a research collaborative has reported new insights into the electronic conduction and interference on 2-D topological insulators—an exotic kind of insulator that ...

Recommended for you



Researchers develop interesting building blocks for quantum computers 6 September 13, 2018

Researchers at the MESA+ research institute of the University of Twente, working together with colleagues in Delft and Eindhoven, have successfully developed nanowires allowing


individual electrons to be captured by a 'quantum ...



Quantum simulation reveals mobility edge in a low-dimensional disordered landscape

August 21, 2018

A recent flood of research scholarship on electronic transport in low-dimensional (2-D or 1D) materials like graphene or carbon nanotubes reflects the tremendous potential of these materials to unveil a deeper understanding ...



An ultracompact laser has been made using nanoscale semiconductor columns

September 13, 2018

A tiny laser comprising an array of nanoscale semiconductor cylinders (see image) has been made by an all-A*STAR team. This is the first time that lasing has been achieved in non-metallic nanostructures, and it promises to ...



Novel insulators with conducting

User comments



Please sign in to add a comment. Registration is free, and takes less than a minute. [Read more](#)

[Sign in](#)

[Click here to reset your password.](#)

[Sign in to get notified via email when new comments are made.](#)

We recommend

[Bilinear magnetoelectric resistance as a probe of three-dimensional spin texture in topological surface states](#)

Nature Physics

[Ballistic Majorana nanowire devices](#)

Nature Nanotechnology

[Numerical analysis of band tails in nanowires and their effects on the performance of tunneling field-effect transistors](#)

Takahisa Tanaka et al., Japanese Journal of Applied Physics

[Fermi surface in the absence of a Fermi liquid in the Kondo insulator SmB6](#)

Nature Physics

[The quantum Hall effect](#)

David K Ferry et al., Transport in Semiconductor Mesoscopic Devices

[Current-driven magnetic domain wall motion and its real-time detection](#)

Kab-Jin Kim et al., Japanese Journal of Applied Physics

Powered by **TREND MD**



- Top
- Home
- Search
- Mobile version
- Help
- FAQ
- About
- Contact
- Science X Account
- Sponsored Account
- Newsletter
- RSS feeds
- Feature Stories
- Latest news
- Week's top
- Archive
- Android app
- iOS app
- Amazon Kindle

Connect

