

Geometry of an electron determined for the first time

Date: May 23, 2019

Source: University of Basel

Summary: Physicists are able to show for the first time how a single electron looks in an artificial atom. A newly developed method enables them to show the probability of an electron being present in a space. This allows improved control of electron spins, which could serve as the smallest information unit in a future quantum computer.

FULL STORY



Quantum computing concept (stock image).

Credit: © metamorworks / Adobe Stock

Physicists at the University of Basel are able to show for the first time how a single electron looks in an artificial atom. A newly developed method enables them to show the probability of an electron being present in a space. This allows improved control of electron spins, which could serve as the smallest information unit in a future quantum computer. The experiments were published in *Physical Review Letters* and the related theory in *Physical Review B*.

The spin of an electron is a promising candidate for use as the smallest information unit (qubit) of a quantum computer. Controlling and switching this spin or coupling it with other spins is a challenge on which numerous research groups worldwide are working. The stability of a single spin and the entanglement of various spins depends, among other things, on the geometry of the electrons -- which previously had been impossible to determine experimentally.

Only possible in artificial atoms

Scientists in the teams headed by professors Dominik Zumbühl and Daniel Loss from the Department of Physics and the Swiss Nanoscience Institute at the University of Basel have now developed a method by which they can spatially determine the geometry of electrons in quantum dots.

A quantum dot is a potential trap which allows to confine free electrons in an area which is about 1000 times larger than a natural atom. Because the trapped electrons behave similar to electrons bound to an atom, quantum dots are also known as "artificial atoms."

The electron is held in the quantum dot by electric fields. However, it moves within the space and, with different probabilities corresponding to a wave function, remains in certain locations within its confinement.

Charge distribution sheds light

The scientists use spectroscopic measurements to determine the energy levels in the quantum dot and study the behavior of these levels in magnetic fields of varying strength and orientation. Based on their theoretical model, it is possible to determine the electron's probability density and thus its wave function with a precision on the sub-nanometer scale.

"To put it simply, we can use this method to show what an electron looks like for the first time," explains Loss.

Better understanding and optimization

The researchers, who work closely with colleagues in Japan, Slovakia and the US, thus gain a better understanding of the correlation between the geometry of electrons and the electron spin, which should be stable for as long as possible and quickly switchable for use as a qubit.

"We are able to not only map the shape and orientation of the electron, but also control the wave function according to the configuration of the applied electric fields. This gives us the opportunity to optimize control of the spins in a very targeted manner," says Zumbühl.

The spatial orientation of the electrons also plays a role in the entanglement of several spins. Similarly to the binding of two atoms to a molecule, the wave functions of two electrons must lie on one plane for successful entanglement.

With the aid of the developed method, numerous earlier studies can be better understood and the performance of spin qubits can be further optimized in the future.

Story Source:

Materials provided by **University of Basel**. *Note: Content may be edited for style and length.*

Related Multimedia:

- Illustration of an electron is trapped in a quantum dot
-

Journal References:

1. Leon C. Camenzind, Liuqi Yu, Peter Stano, Jeremy D. Zimmerman, Arthur C. Gossard, Daniel Loss, Dominik M. Zumbühl. **Spectroscopy of Quantum Dot Orbitals with In-Plane Magnetic Fields**. *Physical Review Letters*, 2019; 122 (20) DOI: 10.1103/PhysRevLett.122.207701
 2. Peter Stano, Chen-Hsuan Hsu, Leon C. Camenzind, Liuqi Yu, Dominik Zumbühl, Daniel Loss. **Orbital effects of a strong in-plane magnetic field on a gate-defined quantum dot**. *Physical Review B*, 2019; 99 (8) DOI: 10.1103/PhysRevB.99.085308
-

Cite This Page:

University of Basel. "Geometry of an electron determined for the first time." ScienceDaily. ScienceDaily, 23 May 2019. <www.sciencedaily.com/releases/2019/05/190523104942.htm>.

RELATED STORIES

Long-Distance Quantum Information Exchange -- Success at the Nanoscale

Mar. 18, 2019 — Researchers have realized the swap of electron spins between distant quantum dots. The discovery is a step towards applications of quantum information, as the dots leave enough room for delicate ... [read more »](#)

Two Quantum Dots Are Better Than One: Using One Dot to Sense Changes in Another

Sep. 19, 2018 — Researchers developed the first device that can detect single-electron events in a self-assembled quantum dot in real time. The device detects the single-electron tunneling events of one quantum dot ... [read more »](#)

Ultra-Bright, Highly Coherent Electron Source Developed to Improve the Performance of Electron-Beam Analytical Instrument

Jan. 28, 2016 — A new technology has been developed to fabricate lanthanum hexaboride (LaB6) single-crystal nanowires with clean surface, a promising cold field-emission electron source for electron microscopes and ... [read more »](#)

Improved Stability of Electron Spins in Qubits

Sep. 7, 2015 — Calculation with electron spins in a quantum computer assumes that the spin states last for a sufficient period of time. Physicists have now demonstrated that electron exchange in quantum dots ... [read more »](#)